

Clinical Policy: Rolapitant (Varubi)

Reference Number: CP.PMN.102

Effective Date: 02.01.17

Last Review Date: 08.25

Line of Business: HIM, Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Rolapitant (Varubi™) is a substance P/neurokinin 1 (NK₁) receptor antagonist.

FDA Approved Indication(s)

Varubi is indicated in combination with other antiemetic agents in adults for the prevention of delayed nausea and vomiting associated with initial and repeat courses of emetogenic cancer chemotherapy, including, but not limited to, highly emetogenic chemotherapy.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Varubi is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Prevention of Nausea and Vomiting Associated with Cancer Chemotherapy (must meet all):**

1. Prescribed for the prevention of chemotherapy-induced nausea/vomiting;
2. Age ≥ 18 years;
3. Member is scheduled to receive moderately to highly emetogenic cancer chemotherapy (*see Appendix D*);
4. Member meets one of the following (a or b):*
**For Illinois HIM requests, the step therapy requirements below do not apply as of 1/1/2026 per IL HB 5395.*
 - a. Failure of aprepitant, unless contraindicated or clinically significant adverse effects are experienced;
**Prior authorization may be required for aprepitant*
 - b. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
5. Prescribed in combination with a serotonin (5-HT₃) receptor antagonist (*ondansetron is preferred*) and dexamethasone;
6. Dose does not exceed both of the following (a and b):
 - a. 180 mg every 2 weeks;
 - b. 2 tablets every 2 weeks.

Approval duration: Projected duration of chemotherapy

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace and CP.PMN.16 for Medicaid; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Prevention of Nausea and Vomiting Associated with Cancer Chemotherapy (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy;
3. Member continues to receive moderately to highly emetogenic cancer chemotherapy (*see Appendix D*);
4. Prescribed in combination with a 5-HT₃ receptor antagonist (*ondansetron is preferred*) and dexamethasone;
5. If request is for a dose increase, new dose does not exceed both of the following (a and b):
 - a. 180 mg every 2 weeks;
 - b. 2 tablets every 2 weeks.

Approval duration: Projected duration of chemotherapy

B. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: HIM.PA.33 for health insurance marketplace and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: HIM.PA.103 for health insurance marketplace and CP.PMN.16 for Medicaid; or

2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid, or evidence of coverage documents

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

5-HT₃: serotonin 5-hydroxytryptamine, type 3

FDA: Food and Drug Administration

NCCN: National Comprehensive Cancer Network

NK₁: neurokinin 1

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
aprepitant (Emend®)	125 mg PO on day 1 and then 80 mg PO on days 2 and 3 of each chemotherapy cycle	Per chemotherapy cycle: Day 1: 125 mg Days 2 and 3: 80 mg

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s):
 - CYP2D6 substrates with a narrow therapeutic index (e.g., thioridazine and pimozide)
 - Pediatric patients less than 2 years of age due to irreversible impairment of sexual development and fertility in juvenile rats
- Boxed warning(s): none reported

Appendix D: American Society of Clinical Oncology (ASCO) and National Comprehensive Cancer Network (NCCN) Recommendations in Oncology

- Minimal emetic risk chemotherapy: No routine prophylaxis is recommended.
- Low emetic risk chemotherapy: Recommended options include dexamethasone (recommended by both ASCO and NCCN) or metoclopramide, prochlorperazine, or a 5-HT₃ receptor antagonist (recommended by NCCN only). NK₁ receptor antagonists are not included in low risk antiemetic recommendations.

- Moderate emetic risk chemotherapy: 5-HT₃ receptor antagonists and dexamethasone may be used in combination and with or without NK₁ receptor antagonists. Olanzapine may also be used in combination with palonosetron and dexamethasone.
 - Examples of moderate emetic risk chemotherapy: bendamustine, carboplatin, clofarabine, cyclophosphamide $\leq 1,500$ mg/m², cytarabine > 200 mg/m², daunorubicin, doxorubicin < 60 mg/m², epirubicin ≤ 90 mg/m², idarubicin, ifosfamide, irinotecan, oxaliplatin
- High emetic risk chemotherapy: NK₁ receptor antagonists are recommended for use in combination with 5-HT₃ receptor antagonists and dexamethasone. Olanzapine may also be used in combination with 5-HT₃ receptor antagonists, dexamethasone, and/or NK₁ receptor antagonists.
 - Examples of high emetic risk chemotherapy: carmustine, cisplatin, cyclophosphamide $> 1,500$ mg/m², dacarbazine, mechlorethamine, streptozocin, fam-trastuzumab deruxtecan-nxki
- Breakthrough emesis: Per NCCN, an agent from a different drug class is recommended to be added to the current antiemetic regimen. Drug classes include atypical antipsychotics (olanzapine), benzodiazepines (lorazepam), cannabinoids (dronabinol), phenothiazines (prochlorperazine, promethazine), 5-HT₃ receptor antagonists (dolasetron, ondansetron, granisetron), steroids (dexamethasone), or haloperidol, metoclopramide, scopolamine. An NK₁ receptor antagonist may be added to the prophylaxis regimen of the next chemotherapy cycle if not previously included.

Appendix E: States with Regulations against Redirections in Stage IV or Metastatic Cancer

State	Step Therapy Prohibited?	Notes
FL	Yes	For stage 4 metastatic cancer and associated conditions.
GA	Yes	For stage 4 metastatic cancer. Redirection does not refer to review of medical necessity or clinical appropriateness.
IA	Yes	For standard of care stage 4 cancer drug use, supported by peer-reviewed, evidence-based literature, and approved by FDA.
LA	Yes [‡]	For stage 4 advanced, metastatic cancer or associated conditions. [‡] Exception if clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy.
MS	Yes	<i>*Applies to HIM requests only*</i> For advanced metastatic cancer and associated conditions
NV	Yes	Stage 3 and stage 4 cancer patients for a prescription drug to treat the cancer or any symptom thereof of the covered person
OH	Yes	<i>*Applies to Commercial and HIM requests only*</i> For stage 4 metastatic cancer and associated conditions
OK	Yes	<i>*Applies to HIM requests only*</i> For advanced metastatic cancer and associated conditions
PA	Yes	For stage 4 advanced, metastatic cancer
TN	Yes [^]	For stage 4 advanced metastatic cancer, metastatic blood cancer, and associated conditions

State	Step Therapy Prohibited?	Notes
		^Exception if step therapy is for AB-rated generic equivalent, interchangeable biological product, or biosimilar product to the equivalent brand drug
TX	Yes	For stage 4 advanced, metastatic cancer and associated conditions

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Prevention of chemotherapy-induced nausea and vomiting	180 mg as a single dose 2 hours prior to the initiation of each chemotherapy, but at no less than 2 week intervals. <i>Administer in combination with dexamethasone and a 5-HT₃ receptor antagonist</i>	180 mg

VI. Product Availability

Tablet: 90 mg

VII. References

1. Varubi Prescribing Information. Waltham, MA: Tesaro, Inc.; August 2020. Available at: https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/206500s008,208399s0041bl.pdf. Accessed April 18, 2025.
2. Hesketh, PJ, Kris MG, Basch E, et al. Antiemetics: American Society of Clinical Oncology Guideline Update. *J Clin Oncol*. 2020. 38:2,782-2,797. doi.org/10.1200/JCO.20.01296.
3. National Comprehensive Cancer Network. Antiemesis Version 1.2025. Available at https://www.nccn.org/professionals/physician_gls/pdf/antiemesis.pdf. Accessed May 12, 2025.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J8670	Rolapitant, oral, 1 mg
J2797	Injection, Rolapitant, 0.5 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2021 annual review: no significant changes; updated contraindications and dosing information per PI; references to	11.13.20	02.21

Reviews, Revisions, and Approvals	Date	P&T Approval Date
HIM.PHAR.21 revised to HIM.PA.154; updated HCPCS code; references reviewed and updated.		
Added allowance for bypassing redirection if state regulations do not allow step therapy in Stage IV or metastatic cancer settings with additional details in appendix E.	04.27.21	
Added Nevada to Appendix E.	08.03.21	
1Q 2022 annual review: no significant changes; removed IV formulation as product is no longer on the market; references reviewed and updated.	10.04.21	02.22
Template changes applied to other diagnoses/indications and continued therapy section.	09.26.22	
1Q 2023 annual review: no significant changes; modified to generalize beyond Stage IV or metastatic cancer to the following redirection bypass: “Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings”; references reviewed and updated.	10.10.22	02.23
3Q 2023 annual review: no significant changes; clarified quantity and dose limit as separate requirements; references reviewed and updated; updated Appendix E to include Oklahoma.	04.19.23	08.23
Updated Appendix E to include Mississippi.	06.05.24	
3Q 2024 annual review: no significant changes; added HCPCS code J2797; references reviewed and updated.	05.06.24	08.24
3Q 2025 annual review: no significant changes; added step therapy bypass for IL HIM per IL HB 5395; updated Appendix E with revised language and exception for Tennessee; references reviewed and updated.	04.18.25	08.25

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering

benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.