

We value everything you do to deliver quality care to our members – your patients – and to ensure they have a positive healthcare experience. There are several HEDIS[®] behavioral health measures that providers can directly impact related to follow-up care for mental illness or substance use disorders, medication adherence and metabolic monitoring. That’s why we’ve created this easy-to-use At-A-Glance Toolkit. It will give you the tools you need to meet, document and code HEDIS[®] measures. Together, we can provide the care and services our members need to stay healthy while also improving our quality scores and Star Ratings. Please contact your Provider Relations representative if you need more information or have any questions.

Quality care is a team effort. Thank you for playing a starring role!

HEDIS’s technical specifications may not align directly with some state guidelines or access standards. Providers are expected to meet state standards.

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(ADD-E) Follow-up care for Children Prescribed ADHD Medication ●</p> <p>At least three follow-up visits within 300-day (10-month) period, one of which was within 30 days of when the first ADHD medication was dispensed.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> • Initiation Phase. • Continuation and Maintenance (C&M) Phase. 	<p>Tips:</p> <ul style="list-style-type: none"> • Complete a medical and psychiatric examination prior to diagnosing and prescribing ADHD medications. • Limit the first prescription of ADHD medication to a 14-to-21-day supply. • Have member or parent schedule a follow-up appointment before leaving the office when a new ADHD medication has been prescribed. 	<p>Outpatient Visit: CPT: 90797,90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255</p> <p>POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72</p> <p>BH Outpatient Visit: CPT: 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99510, 99483, 99492-99494</p> <p>HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013, H2014, H2015, H2016, H2017, H2018, H2019, H2020, T1015</p> <p>Observation Visit: CPT: 99217-99220</p> <p>Health and Behavior Assessment/Intervention: CPT: 96150-96154, 96156, 96158, 96159, 96164, 96165, 96167, 96168, 96170, 96171</p> <p>Visit Setting Unspecified Value Set with Partial Hospitalization POS: CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255.</p> <p>POS: 52</p> <p>Partial Hospitalization/ Intensive Outpatient: HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485</p> <p>Telehealth Visit: CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255</p> <p>POS: 02, 10</p> <p>Telephone Visits: CPT: 98966-98968, 99441-99443</p> <p>E-visit/Virtual Check-In: CPT: 98969-98972, 99421-99423, 99444, 99457</p> <p>HCPCS: G0071, G2010, G2012, G2061, G2062, G2063</p> <p>Visit Setting Unspecified Value Set with Community Mental Health Center POS*: CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255</p> <p>POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11)</p> <p>Narcolepsy: ICD-10: G47.411, G47.419, G47.421, G47.429</p>



Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(AMM) Antidepressant Medication Management ● ● ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Diagnosed with Major Depression and remain on antidepressant medication treatment.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> Acute: at least 12 weeks and Continuation: at least 6 months. 	<p>Tips</p> <ul style="list-style-type: none"> Educate patients on the importance of taking antidepressants Monitor response to treatment with a standardized tool such as the Patient Healthcare Questionnaire (PHQ-9). Discuss the 988 Suicide & Crisis Lifeline (988lifeline.org) with patients and family 	<p>Antidepressant Medications</p> <ul style="list-style-type: none"> Miscellaneous Antidepressants: Bupropion, Vilazodone, Vortioxetine Monoamine Oxidase Inhibitors: Isocarboxazid, Selegiline, Phenelzine, Tranylcypromine Phenylpiperazine Antidepressants: Nefazodone, Trazodone Psychotherapeutic Combinations: Amitriptyline-chlordiazepoxide, Fluoxetine-olanzapine, Amitriptyline-perphenazine SNRI Antidepressants: Desvenlafaxine, Duloxetine, Venlafaxine, Levomilnacipran SSRI Antidepressants: Citalopram, Escitalopram, Fluoxetine, Fluvoxamine, Paroxetine, Sertraline Tetracyclic Antidepressants: Maprotiline, Mirtazapine Tricyclic Antidepressants: Amitriptyline, Amoxapine, Clomipramine, Desipramine, Doxepin (>6 mg), Imipramine, Nortriptyline, Protriptyline, Trimipramine
<p>(APM-E) Metabolic Monitoring for Children and Adolescents on Antipsychotics ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Children and adolescents (1-17 years of age) who had 2 or more antipsychotic prescriptions and had metabolic testing.</p> <p>Three Rates:</p> <ul style="list-style-type: none"> Blood glucose testing Cholesterol testing Both blood glucose and cholesterol testing 	<p>Tips</p> <ul style="list-style-type: none"> Provide members/caregivers with lab orders for HbA1c or glucose and cholesterol or LDL-C to be completed yearly. Educate the member and caregiver about the risks associated with taking antipsychotic medications and the importance of regular follow up care. Submit applicable codes. 	<p>Description (Need either A1c or Glucose and LCL-C or Cholesterol) & Codes*</p> <ul style="list-style-type: none"> HbA1c Lab Tests: <ul style="list-style-type: none"> CPT: 83036, 83037 CPT-CAT-II: 3044F, 3046F, 3051F, 3052F Glucose Lab Tests <ul style="list-style-type: none"> CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951 LDL-C Lab Tests <ul style="list-style-type: none"> CPT: 80061, 83700, 83701, 83704, 83721 CPT-CAT-II: 3048F, 3049F, 3050F Cholesterol Lab Tests <ul style="list-style-type: none"> CPT: 82465, 83718, 83722, 84478 <p><i>Note: Do not include a modifier when using CPT-CAT-II codes.</i></p>
<p>(APP) Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Children and adolescents (1-17 years of age) had documentation of psychosocial care as first line treatment prior to antipsychotic medication prescription.</p>	<p>Tips</p> <ul style="list-style-type: none"> Before prescribing antipsychotic medication, complete or refer for a trial of first-line psychosocial care. Antipsychotic medications should be a part of a comprehensive, multi-modal plan for coordinated treatment that includes psychosocial care. Periodically the ongoing need for continued therapy with antipsychotic medication should be reviewed. 	<p>Description & Codes*</p> <ul style="list-style-type: none"> Psychosocial Care <ul style="list-style-type: none"> CPT: 90832-90839, 90840-90849, 90853, 90875, 90876, 90880 HCPCS: G0176, G0177, G0409-G0411, H0004, H0035-H0039, H0040, H2000-H2020, S0201, S9480-S9485 Residential Behavioral Health Treatment <ul style="list-style-type: none"> HCPCS: H0017-H0019, T2048

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services			
(COU) Risk of Continued Opioid Use ● ● Members 18 years of age and older at risk for continued opioid use: <ul style="list-style-type: none"> 15 days of prescription within 30-day period 31 days of prescription within 62-day period Note: lower rate indicates better performance.	Tips <ul style="list-style-type: none"> Only prescribe opioids when medically necessary, in the lowest effective dose, for the shortest duration necessary. Assess the risks and benefits with patients within one to four weeks of initiating opioid therapy for chronic pain. Educate on the risks of long-term opioid use. 	Opioid Medications <ul style="list-style-type: none"> Benzhydrocodone, Buprenorphine (transdermal patch and buccal film), Butorphanol, Codeine, Dihydrocodeine, Fentanyl, Hydrocodone, Hydromorphone, Levorphanol, Meperidine, Methadone, Morphine, Opium, Oxycodone, Pentazocine, Tapentadol, Tramadol The following opioid medications are excluded from this measure: <ul style="list-style-type: none"> Injectables Opioid-containing cough and cold products. Single-agent and combination buprenorphine products used as part of medication-assisted treatment of opioid use disorder (buprenorphine sublingual tablets, buprenorphine subcutaneous implant and all buprenorphine/naloxone combination products). Ionsys[®] (fentanyl transdermal patch). 			
(DSF-E) Depression Screening and Follow-Up for Adolescents and Adults ● ● ● ● 12 years of age and older screened for clinical depression using a standardized instrument and received follow-up care if screened positive. Two Rates: <ul style="list-style-type: none"> Depression Screening Follow-up on Positive screen 	Tips <ul style="list-style-type: none"> Use age-appropriate screening instruments. Train staff on the importance of depression screenings and to recognize the risk factors for depression. Work with a care team to coordinate follow-up care for members with a positive screening. Ensure all services conducted during the visit are coded appropriately, including the depression screening LOINC codes. Coordinate file submissions to the health plan that include HER data. 	Instruments for Adolescents (≤ 17 years)			
		Patient Health Questionnaire (PHQ-9) [®]		Total Score LOINC Codes	Positive Finding
		Patient Health Questionnaire Modified for Teens (PHQ-9M) [®]		44261-6	Total Score ≥ 10
		Patient Health Questionnaire-2 (PHQ-2) [®]		89204-2	Total Score ≥ 10
		Beck Depression Inventory – Fast Screen (BDI-FS) [®]		55758-7	Total score ≥ 3
		Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)		89208-3	Total score ≥ 8
		Edinburgh Postnatal Depression Scale (EPDS)		89205-9	Total score ≥ 17
		PROMIS Depression		99046-5	Total score ≥ 10
				71965-8	Total score (T Score) ≥ 60
		Instruments for Adults (18 + years)			
		Patient Health Questionnaire (PHQ-9) [®]		Total Score LOINC Codes	Positive Finding
		Patient Health Questionnaire-2 (PHQ-2) [®]		44261-6	Total Score ≥ 10
		Beck Depression Inventory – Fast Screen (BDI-FS) [®]		55758-7	Total score ≥ 3
		Beck Depression Inventory (BDI-II)		89208-3	Total score ≥ 8
		Beck Depression Inventory (BDI-II)		89209-1	Total score ≥ 20
		Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)		89205-9	Total score ≥ 17
		Duke Anxiety – Depression Scale (DUKE AD) [®]		90853-3	Total score ≥ 30
Geriatric Depression Scale Short Form (GDS)		48545-8	Total score ≥ 5		
Geriatric Depression Scale Long Form (GDS)		48544-1	Total score ≥ 10		
Edinburgh Postnatal Depression Scale (EPDS)		99046-5	Total score ≥ 10		
My Mood Monitor (M-3) [®]		71777-7	Total score ≥ 5		
PROMIS Depression		71965-8	Total score (T Score) ≥ 60		
Clinically Useful Depression Outcome Scale (CUDOS)		90221-3	Total score ≥ 31		

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(FUA) Follow-Up After Emergency Department Visit with Substance Use Disorder ● ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Discharges age 13 years and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose for which there is follow up.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> • Within 30 days of discharge • Within 7 days of discharge 	<p>Tips</p> <p>Please follow the KanCare Mental Health access standards as listed in the Sunflower Provider Manual. This will also ensure compliance with FUA.</p> <ul style="list-style-type: none"> • Offer virtual, telehealth, and phone visits. • Maintain appointment availability in your practice for patients and schedule follow-up appointments before the patient leaves the office. • Discuss the benefits of seeing a primary or specialty provider. • Offer mutual help options like case management, peer recovery support, harm reduction, 12-step fellowships (AA, NA, etc.), or other community support groups. • Reach out proactively within 24 hours if the patient does not keep scheduled appointment to schedule another. <p>The visit can be with any practitioner if the claim includes a diagnosis of SUD (e.g., F10.xx-F19.xx) or drug overdose (e.g., T40-T43, T51). If the visit occurs with a mental health provider, the claim does not have to include the SUD or drug overdose diagnosis.</p>	<p>Description & Codes*</p> <p>Outpatient Visit with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492-99494, 99510 • HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72 <p>Intensive Outpatient Encounter or Partial Hospitalization with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx POS: 52 <p>Non-residential Substance Abuse Treatment Facility with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx POS: 57, 58 <p>Community Mental Health Center* Visit with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11) <p>Observation Visit with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 99217, 99218, 99219, 99220 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx <p>Peer Support Service with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • HCPCS: G0177, H0024, H0025, H0038-H0040, H0046, H2014, H2023, S9445, T1012, T1016 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx <p>Opioid Treatment Service That Bills Monthly or Weekly with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • HCPCS: G2086, G2087, G2071, G8074-G2077, G2080 ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx <p>Telehealth Visit with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx POS: 02, 10 <p>Telephone Visit with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 98966-98968, 99441-99443 • ICD-10: F10.xx-F19.xx or T40.xxxx-T43.xxxx, T51.xxxx <p>E-Visit or Virtual Check In with any Diagnosis of SUD or Drug Overdose</p> <ul style="list-style-type: none"> • CPT: 98969-98972, 99421-99444, 99457, 99458 • HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252 <p>Substance Use and Substance Use Disorder Services</p> <ul style="list-style-type: none"> • CPT: 99408, 99409 • HCPCS: T1012, G0396, G0397, H0001, H0005, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, H0006, H0028

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS® Measure	Best Practice	Coding and Services
		<ul style="list-style-type: none"> Behavioral Health Screening or Assessment for SUD or Mental Health Disorders <ul style="list-style-type: none"> CPT: 99408, 99409 HCPCS: G0396, G0397, G0442, H2011, H0001, H0002, H0031, H0049 Pharmacotherapy Dispensing Event or Medication Treatment Event <ul style="list-style-type: none"> Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral; delayed-release tablet), Buprenorphine (implant, injection, or sublingual tablet), Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) HCPCS: G2069, G2070, G2072, G2073, H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109
<p>(FUH) Follow-Up After Hospitalization for Mental Illness ● ● ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Discharges 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and had a follow-up visit with mental health provider.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> Within 30 days after discharge Within 7 days after discharge 	<p>Tips</p> <p>Please follow the KanCare Mental Health access standards as listed in the Sunflower Provider Manual. This will also ensure compliance with FUH.</p> <ul style="list-style-type: none"> Schedule follow up appointments prior to discharge and include the date and time on discharge instructions. Submit applicable codes. Offer telehealth and phone visits. Reach out proactively to assist in (re)scheduling appointments within the required timeframes. Partner with the health plan to address social determinants, health equity, and quality care. 	<p>Description & Codes*</p> <ul style="list-style-type: none"> Outpatient Visit with a Mental Health Provider <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99492-99494, 99510, 99483 POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H0039, H0040, H2000, H2010, H2011, H2013-H2020, T1015 Visit Setting Unspecified Value Set with Partial Hospitalization POS with Mental Health Provider <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 52 Partial Hospitalization/Intensive Outpatient <ul style="list-style-type: none"> HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 Visit Setting Unspecified Value Set with Community Mental Health Center POS* <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11) Electroconvulsive Therapy with Ambulatory Surgical Center POS/ Community Mental Health Center POS*/Outpatient POS/Partial Hospitalization POS <ul style="list-style-type: none"> CPT: 90870 Ambulatory POS: 24 Comm. POS: 53 *Certified Community Behavioral Health Clinics POS: 11 Partial Hosp. POS: 52 Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72 Telehealth Visit <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 POS: 02, 10 Observation <ul style="list-style-type: none"> CPT: 99217-99220 Transitional Care Management <ul style="list-style-type: none"> CPT: 99495, 99496 Telephone Visit <ul style="list-style-type: none"> CPT: 98966-98968, 99441-99443 Psychiatric Collaborative Care Management <ul style="list-style-type: none"> CPT: 99492-99494 HCPCS: G0512

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS® measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(FUI) Follow-Up After High-Intensity Care for Substance Use Disorder</p> <p>● ●</p> <p>Discharges 13 years of age and older from acute inpatient hospitalizations, residential treatment or withdrawal management visits that result in a follow-up visit or service for substance use disorder.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> Follow-up within 30 days of discharge and, Follow-up within 7 days of discharge <p><i>Note: Follow-up does not include withdrawal management.</i></p>	<p>Tips</p> <p>Please follow the KanCare Mental Health access standards as listed in the Sunflower Provider Manual. This will also ensure compliance with FUI.</p> <ul style="list-style-type: none"> Offer virtual, telehealth and phone visits. Maintain appointment availability in your practice for patients and schedule follow-up appointments before the patient leaves the office. Offer mutual help options like case management, peer recovery support, harm reduction, 12-steps fellowships (AA, NS, etc.), or other community support groups. Reach out proactively within 24 hours if the patient does not keep scheduled appointment to schedule another. 	<p>Description & Codes*</p> <ul style="list-style-type: none"> An Acute or Nonacute Inpatient Admission or Residential Behavioral Health Stay with a Principal Diagnosis of SUD on the Discharge Claim (does not include visits that occur on the date of the episode) <ul style="list-style-type: none"> F10.xx-F19.xx Outpatient Visit with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: F10.xx-F19.xx POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72 Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 ICD-10: F10.xx-F19.xx POS: 52 Non-residential Substance Abuse Treatment Facility with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 57, 58 Community Mental Health Center* Visit with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11) Telehealth Visit with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F19.xx POS: 02, 10 Substance use Disorder Services with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 99408, 99409 HCPCS: T1012, G0396, G0397, H0001, H0005, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, H0006, H0028 Opioid Treatment Service that Bills Monthly or Weekly with a Principal Diagnosis of SUD <ul style="list-style-type: none"> HCPCS: G2086, G2087, G2071, G8074-G2077, G2080 ICD-10: F10.xx-F19.xx Observation Visit with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 99217, 99218, 99219, 99220 ICD-10: F10.xx-F19.xx Residential Behavioral Health Treatment with a Principal Diagnosis of SUD <ul style="list-style-type: none"> HCPCS: H0017, H0018, H0019, T2048 ICD-10: F10.xx-F19.xx Telephone Visit with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 98966-98968, 99441-99443 ICD-10: F10.xx-F19.xx

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS® Measure	Best Practice	Coding and Services
		<ul style="list-style-type: none"> E-Visit or Virtual Check in with a Principal Diagnosis of SUD <ul style="list-style-type: none"> CPT: 98969-98972, 99421-99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252 Pharmacotherapy Dispensing Event or Medication Treatment Event <ul style="list-style-type: none"> Medications: Disulfiram (oral), Naltrexone (oral and injectable), Acamprosate (oral; delayed release tablet), Buprenorphine (implant, injection, or sublingual tablet), Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) HCPCS: G2069, G2070, G2072, G2073, H0020, H0033, J0570-J0575, J2315, Q9991, Q9992, S0109
<p>(FUM) Follow-Up After Emergency Department Visit for Mental Illness</p> <p>● ●</p> <p><i>*Applicable Foster Care Measure</i></p> <p>Emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> Follow-up within 30 days of the ED visit (31 total days). Follow-up within 7 days of the ED visit (8 total days). 	<p>Tips</p> <p>Please follow the KanCare Mental Health access standards as listed in the Sunflower Provider Manual. This will also ensure compliance with FUM.</p> <ul style="list-style-type: none"> Offer virtual, telehealth and phone visits. Maintain appointment availability in your practice for patients and schedule follow-up appointments before the patient leaves the office. Discuss benefits of seeing a primary or specialty provider and appropriate ED utilization. Partner with the health plan to address social determinants, health equity, and quality care. 	<p>Description & Codes*</p> <ul style="list-style-type: none"> Outpatient Visit with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492 HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 ICD-10: F10.xx-F99 POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72 Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 ICD-10: F10.xx-F99 POS: 52 Community Mental Health Center* Visit with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F99 POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11) Electroconvulsive therapy with an Outpatient POS and with a principal diagnosis of intentional self-harm and with any diagnosis of a mental health disorder <ul style="list-style-type: none"> CPT: 90780 ICD-10: T40.xxxx-T43.xxxx, T51.xxxx POS: 03, 05, 07, 09, 11-20, 22, 24, 33, 49, 50, 52, 53, 71, 72 Telehealth Visit with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 ICD-10: F10.xx-F99 POS: 02, 10 Observation Visit with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 99217, 99218, 99219, 99220 ICD-10: F10.xx-F99 Telephone Visit with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 98966-98968, 99441-99443 ICD-10: F10.xx-F99 E-Visit or Virtual Check in with a Principal Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> CPT: 98969-98972, 99421-99444, 99457, 99458 HCPCS: G0071, G2010, G2012, G2061-G2063, G2250-G2252

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS® measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
		<ul style="list-style-type: none"> • An Outpatient with a Principal Diagnosis of Intentional Self-Harm with a Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255, 98960-98962, 99078, 99201-99205, 99211-99215, 99241-99245, 99341-99345, 99347-99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99483, 99492 • HCPCS: G0155, G0176, G0177, G0409, G0463, G0512, H0002, H0004, H0031, H0034, H0036, H0037, H2000, H2010, H2013, H2015, H2017-H2020, T1015 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 • POS: 03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71-72 • Intensive Outpatient Encounter or Partial Hospitalization with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • HCPCS: G0410, G0411, H0035, H2001, H2012, S0201, S9480, S9484, S9485 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 POS: 52 • Community Mental Health Center* Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 • POS: 53 (*Certified Community Behavioral Health Clinics: POS: 11) • Electroconvulsive Therapy with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 90780 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 POS: 03, 05, 07, 09, 11-20, 22, 24, 33, 49, 50, 52, 53, 71, 72 • Telehealth Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 90791, 90792, 90832, 90833, 90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221-99223, 99231-99233, 99238, 99239, 99251-99255 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 • POS: 02, 10 • Observation Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 99217, 99218, 99219, 99220 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99 • Telephone Visit with a Principal Diagnosis of Intentional Self-harm with any Diagnosis of a Mental Health Disorder <ul style="list-style-type: none"> • CPT: 98966-98968, 99441-99443 • ICD-10: T40.xxxx-T43.xxxx, T51.xxxx with F10.xx-F99

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(IET) Initiation and Engagement of Substance Use Disorder Treatment</p> <p>● ● ●</p> <p>Adolescent and adult members with a new episode of substance use disorder (SUD) episodes that result in treatment initiation and engagement.</p> <ul style="list-style-type: none"> Initiation of SUD treatment: new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter or partial hospitalization, telehealth, or medication treatment within 14 days. Engagement of SUD treatment: new SUD episodes that have evidence of treatment. 	<p>Tips</p> <ul style="list-style-type: none"> Explain the importance of a follow-up to your patients. Schedule an initial follow-up appointment within 14 days. Reschedule patients as soon as possible who do not keep initial appointments. Use telehealth where appropriate. Offer mutual help options like case management, peer recovery support, harm reduction, 12-step fellowships (AA, NA, etc.), or other community support groups. Maintain appointment availability in your practice for patients and schedule follow-up appointments before the patient leaves the office. Submit applicable codes. 	<p>Alcohol Use Disorder Treatment Medications</p> <ul style="list-style-type: none"> Aldehyde Dehydrogenase Inhibitor – Disulfiram (oral) Antagonist – Naltrexone (oral and injectable) Other – Acamprosafe (oral; delayed-release tablet) <p>Opioid Use Disorder Treatment Medications</p> <ul style="list-style-type: none"> Antagonist – Naltrexone (oral), Naltrexone Oral Medication List Antagonist – Naltrexone (injectable), Naltrexone Injection Medication List Partial Agonist – Buprenorphine (sublingual tablet), Buprenorphine oral Medication List <p>Opioid Use Disorder Treatment Medications</p> <ul style="list-style-type: none"> Partial Agonist – Buprenorphine (injection), Buprenorphine Injection Medication List Partial Agonist – Buprenorphine (implant), Buprenorphine Implant Medication List Partial Agonist – Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film), Buprenorphine Naloxone Medication List <p>Description & Codes*</p> <ul style="list-style-type: none"> Initiation and Engagement/Treatment <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90834, 90836-90840, 90845, 90847, 90849, 90853, 90875, 90876, 98960-98962, 99078, 99201-99205, 99211-99215, 99217-99220, 99221-99223, 99231, 99232, 99233, 99238, 99239, 99241-99245, 99341-99345, 99347-99350, 99251-99255, 99381-99387, 99391-99397, 99401-99404, 99408, 99409, 99411, 99412, 99483, 99492-99494, 99510 HCPCS: G0155, G0176, G0177, G0396, G0397, G0409, G0443, G0463, G2086, G2087, G0512, G2067-G2078, G2080, H0001, H0002, H0004, H0005, H0007, H0015, H0016, H0022, H0031, H0034-H0037, H0039, H0040, H0047, H0050, H2000, H2010, H2011, H2013-H2020, H2035, H2036, S0201, S9480, S9484, S9485, T1006, T1012, T1015 POS: 02, 03, 05, 07, 09, 11-20, 22, 33, 49-50, 52-53 (includes Certified Community Behavioral Health Clinics), 57, 58, 71-72 Telephone Visits <ul style="list-style-type: none"> CPT: 98966-98968, 99441-99443 E-visit/Virtual Check-in <ul style="list-style-type: none"> CPT: 98969-98972, 99421-99423, 99444, 99457 HCPCS: G0071, G2010, G2012, G2061-G2063

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024



HEDIS [®] Measure	Best Practice	Coding and Services																																																						
<p>(PND-E) Prenatal Depression Screening ●</p> <p>Members were screened for clinical depression while pregnant and, if positive, received follow-up care.</p> <p>Two Rates:</p> <ul style="list-style-type: none"> Depression Screening – using standardized instrument. Follow-up on Positive Screen – within 30 days of a positive depression screen finding. 	<p>Tips</p> <ul style="list-style-type: none"> Explain the importance of a follow-up to your patients. Schedule an initial follow-up appointment within 14 days. Reschedule patients as soon as possible who do not keep initial appointments. Use telehealth where appropriate. Offer mutual help options like case management, peer recovery support, harm reduction, 12-step fellowships (AA, NA, etc.), or other community support groups. Maintain appointment availability in your practice for patients and schedule follow-up appointments before the patient leaves the office. Submit applicable codes. 	<p>Instruments for Adolescents (≤17 years)</p> <table border="1"> <tr> <td>Patient Health Questionnaire (PHQ-9)[®]</td> <td>Total score ≥ 10</td> <td>44261-6</td> </tr> <tr> <td>Patient Health Questionnaire Modified for Teens (PHQ-9M)[®]</td> <td>Total score ≥ 10</td> <td>89204-2</td> </tr> <tr> <td>Patient Health Questionnaire-2 (PHQ-2)[®]</td> <td>Total score ≥ 3</td> <td>55758-7</td> </tr> <tr> <td>Beck Depression Inventory – Fast Screen (BDI-FS)[®]</td> <td>Total score ≥ 8</td> <td>89208-3</td> </tr> <tr> <td>Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)</td> <td>Total score ≥ 17</td> <td>89205-9</td> </tr> <tr> <td>Edinburgh Postnatal Depression Scale (EPDS)</td> <td>Total score ≥ 10</td> <td>99046-5</td> </tr> <tr> <td>PROMIS Depression</td> <td>Total score (T Score) ≥ 60</td> <td>71965-8</td> </tr> </table>	Patient Health Questionnaire (PHQ-9) [®]	Total score ≥ 10	44261-6	Patient Health Questionnaire Modified for Teens (PHQ-9M) [®]	Total score ≥ 10	89204-2	Patient Health Questionnaire-2 (PHQ-2) [®]	Total score ≥ 3	55758-7	Beck Depression Inventory – Fast Screen (BDI-FS) [®]	Total score ≥ 8	89208-3	Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)	Total score ≥ 17	89205-9	Edinburgh Postnatal Depression Scale (EPDS)	Total score ≥ 10	99046-5	PROMIS Depression	Total score (T Score) ≥ 60	71965-8	<p>Positive Finding</p> <p>LOINC Code (Required for numerator 1)</p>	<p>Instruments for Adults (18 + years)</p> <table border="1"> <tr> <td>Patient Health Questionnaire (PHQ-9)[®]</td> <td>Total score ≥ 10</td> <td>44261-6</td> </tr> <tr> <td>Patient Health Questionnaire-2 (PHQ-2)[®]</td> <td>Total score ≥ 3</td> <td>55758-7</td> </tr> <tr> <td>Beck Depression Inventory – Fast Screen (BDI-FS)[®]</td> <td>Total score ≥ 8</td> <td>89208-3</td> </tr> <tr> <td>Beck Depression Inventory (BDI-II)</td> <td>Total score ≥ 20</td> <td>89209-1</td> </tr> <tr> <td>Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)</td> <td>Total score ≥ 17</td> <td>89205-9</td> </tr> <tr> <td>Duke Anxiety-Depression Scale (DUKE-AD)[®]</td> <td>Total score ≥ 30</td> <td>90853-3</td> </tr> <tr> <td>Edinburgh Postnatal Depression Scale (EPDS)</td> <td>Total score ≥ 10</td> <td>99046-5</td> </tr> <tr> <td>My Mood Monitor (M-3)[®]</td> <td>Total score ≥ 5</td> <td>71777-7</td> </tr> <tr> <td>PROMIS Depression</td> <td>Total score (T Score) ≥ 60</td> <td>71965-8</td> </tr> <tr> <td>Clinically Useful Depression Outcome Scale (CUDOS)</td> <td>Total score ≥ 31</td> <td>90221-3</td> </tr> </table>	Patient Health Questionnaire (PHQ-9) [®]	Total score ≥ 10	44261-6	Patient Health Questionnaire-2 (PHQ-2) [®]	Total score ≥ 3	55758-7	Beck Depression Inventory – Fast Screen (BDI-FS) [®]	Total score ≥ 8	89208-3	Beck Depression Inventory (BDI-II)	Total score ≥ 20	89209-1	Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)	Total score ≥ 17	89205-9	Duke Anxiety-Depression Scale (DUKE-AD) [®]	Total score ≥ 30	90853-3	Edinburgh Postnatal Depression Scale (EPDS)	Total score ≥ 10	99046-5	My Mood Monitor (M-3) [®]	Total score ≥ 5	71777-7	PROMIS Depression	Total score (T Score) ≥ 60	71965-8	Clinically Useful Depression Outcome Scale (CUDOS)	Total score ≥ 31	90221-3	<p>Positive Finding</p> <p>LOINC Code (Required for numerator 1)</p>
Patient Health Questionnaire (PHQ-9) [®]	Total score ≥ 10	44261-6																																																						
Patient Health Questionnaire Modified for Teens (PHQ-9M) [®]	Total score ≥ 10	89204-2																																																						
Patient Health Questionnaire-2 (PHQ-2) [®]	Total score ≥ 3	55758-7																																																						
Beck Depression Inventory – Fast Screen (BDI-FS) [®]	Total score ≥ 8	89208-3																																																						
Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)	Total score ≥ 17	89205-9																																																						
Edinburgh Postnatal Depression Scale (EPDS)	Total score ≥ 10	99046-5																																																						
PROMIS Depression	Total score (T Score) ≥ 60	71965-8																																																						
Patient Health Questionnaire (PHQ-9) [®]	Total score ≥ 10	44261-6																																																						
Patient Health Questionnaire-2 (PHQ-2) [®]	Total score ≥ 3	55758-7																																																						
Beck Depression Inventory – Fast Screen (BDI-FS) [®]	Total score ≥ 8	89208-3																																																						
Beck Depression Inventory (BDI-II)	Total score ≥ 20	89209-1																																																						
Center for Epidemiologic Studies Depression Scale – Revised (CESD-R)	Total score ≥ 17	89205-9																																																						
Duke Anxiety-Depression Scale (DUKE-AD) [®]	Total score ≥ 30	90853-3																																																						
Edinburgh Postnatal Depression Scale (EPDS)	Total score ≥ 10	99046-5																																																						
My Mood Monitor (M-3) [®]	Total score ≥ 5	71777-7																																																						
PROMIS Depression	Total score (T Score) ≥ 60	71965-8																																																						
Clinically Useful Depression Outcome Scale (CUDOS)	Total score ≥ 31	90221-3																																																						
		<p>Description & Codes*</p> <ul style="list-style-type: none"> Perinatal <ul style="list-style-type: none"> ICD-10: Z3A.37, Z3A.38, Z3A.39, Z3A.40, Z3A.41, Z3A.42, Z3A.49, Z3A.01, Z3A.08, Z3A.09, Z3A.10, Z3A.11, Z3A.12, Z3A.13, Z3A.14, Z3A.15, Z3A.16, Z3A.17, Z3A.18, Z3A.19, Z3A.20, Z3A.21, Z3A.22, Z3A.23, Z3A.24, Z3A.25, Z3A.26, Z3A.27, Z3A.28, Z3A.29, Z3A.30, Z3A.31, Z3A.32, Z3A.33, Z3A.34, Z3A.35, Z3A.36 CPT: 59400, 59409, 59410, 59510, 59514, 59515, 59610, 59612, 59614, 59618, 59620, 59622 Behavioral Health Encounter <ul style="list-style-type: none"> CPT: 90791, 90792, 90832-90839, 90845-90849, 90853, 90865-90869, 90870, 90875, 90876, 90880, 90887, 99484, 99492, 99493 Depression Case Management Encounter <ul style="list-style-type: none"> CPT: 99366, 99492-99494 HCPCS: G0512, T1016, T1017, T2022, T2023 																																																						

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS® Measure	Best Practice	Coding and Services
		<ul style="list-style-type: none"> Depression or Other Behavioral Health Condition <ul style="list-style-type: none"> ICD-10: F01.51, F01.511, F01.518, F06.4, F10.180, F10.280, F10.980, F11.188, F11.288, F11.988, F12.180, F12.280, F12.980, F13.180, F13.280, F13.980, F14.180, F14.280, F14.980, F15.180, F15.280, F15.980, F16.180, F16.280, F16.980, F18.180, F18.280, F18.980, F19.180, F19.280, F19.980, F20.0-F20.5, F20.81, F20.89, F20.9, F21-F24, F25.0-F25.9, F28, F29, F30.10-F30.13, F30.2-F30.9, F31.0, F31.10-F31.13, F31.2, F31.30-F31.32, F31.4, F31.5, F31.60-F31.64, F31.70-F31.78, F31.81, F31.89, F31.9, F32.0-F32.9, F32.A, F33.0-F33.9, F34.0-F34.9, F39, F40.00-F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230-F40.233, F40.240-F40.248, F40.290-F40.298, F40.8, F40.9, F41.0-F41.9, F42-F42.9, F43.0, F43.10-F43.12, F43.20-F43.29, F43.8-F43.89, F43.9, F44.89, F45.21, F51.5, F53-F53.1, F60.0-F60.9, F63.0-F63.9, F68.10-F68.13, F68.8, F68.A, F84.0-F84.9, F90.0-F90.9, F91.0-F91.9, F93.0-F93.9, F94.0-F94.9, O90.6, O99.340-O99.345 Follow Up Visit <ul style="list-style-type: none"> CPT: 98960-98968, 98970-98972, 98980, 98981, 99078, 99202-99205, 99211-99215, 99242-99245, 99341-99349, 99350, 99381-99387, 99391-99397, 99401-99404, 99411, 99412, 99421-99423, 99441-99443, 99457, 99458, 99483
(POD) Pharmacotherapy for Opioid Use Disorder  Members 16 years of age and older with a diagnosis of opioid use disorder (OUD) and new OUD pharmacotherapy event; measuring the pharmacotherapy events that lasted at least 180 days.	Tips <ul style="list-style-type: none"> Closely monitor medication prescriptions and do not allow any gap in treatment of 8 or more consecutive days. Offer mutual help like peer recovery support, harm reduction, 12-step fellowships (AA, NA, etc.) Provide timely submission of claims with correct medication name, dosage, frequency, and days covered. 	Description & Codes* <ul style="list-style-type: none"> Opioid Use Disorder (OUD) <ul style="list-style-type: none"> F11.10, F11.120-122, F11.129, F11.13-14, F11.150-151, F11.159, F11.181-182, F11.188, F11.19-20, F11.220-222, F11.229, F11.23-24, F11.250-251, F11.259, F11.281-282, F11.288, F11.29 Antagonist <ul style="list-style-type: none"> Naltrexone (oral or injectable) Partial Agonist <ul style="list-style-type: none"> Buprenorphine (sublingual tablet, injection, or implant) Partial Agonist <ul style="list-style-type: none"> Buprenorphine/ naloxone (sublingual tablet, buccal film, sublingual film) Agonist <ul style="list-style-type: none"> Methadone (oral, medical claim codes H0020, S10109, G2067, G2078)
(SAA) Adherence to Antipsychotic Medication for Individuals with Schizophrenia  Members 18 years of age and older with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for 80% of the year.	Tips <ul style="list-style-type: none"> Consider the use of long-acting injectable antipsychotic medications to increase adherence. Provide education on how to take the medication, expected side effects, and talking to the prescriber before stopping the medication. Submit applicable codes. 	Oral Antipsychotics Aripiprazole, Asenapine, Brexpiprazole, Cariprazine, Clozapine, Haloperidol, Iloperidone, Loxapine, Lumateperone, Lurasidone, Molindone, Olanzapine, Paliperidone, Quetiapine, Risperidone, Ziprasidone, Chlorpromazine, Fluphenazine, Perphenazine, Prochlorperazine, Thioridazine, Trifluoperazine, Amitriptyline-perphenazine, Thiothixene Long-Acting Injections <ul style="list-style-type: none"> Long-acting Injections 14 Days Supply <ul style="list-style-type: none"> Risperidone (excluding Perseris®) Long-acting Injections 28 Days Supply <ul style="list-style-type: none"> Aripiprazole, Aripiprazole lauroxil, Fluphenazine decanoate, Haloperidol decanoate, Olanzapine, Paliperidone palmitate Long-acting Injections 30 Days Supply <ul style="list-style-type: none"> Risperidone (Perseris®)

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS® measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024

HEDIS [®] Measure	Best Practice	Coding and Services
<p>(SSD) Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications ●</p> <p>Members 18-64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test.</p>	<p>Tips</p> <ul style="list-style-type: none"> • Provide members/caregivers with lab orders for HbA1c or glucose and cholesterol or LDL-C to be completed yearly. • Educate the member and caregiver about the risks associated with taking antipsychotic medications and the importance of regular follow up care. • Submit applicable codes. 	<p>Description & Codes*</p> <ul style="list-style-type: none"> • Glucose Lab Test <ul style="list-style-type: none"> • CPT: 80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951 • Glucose Test Result or Finding <ul style="list-style-type: none"> • SNOMED: 166890005 (Blood glucose within reference range), 166891009 (Blood sugar below reference range), 166892002 (Blood sugar above reference range), 442545002 (Blood glucose outside reference range), 444780001 (Glucose above reference range), 1179458001 (Blood glucose below reference range) • HbA1c Lab Test <ul style="list-style-type: none"> • CPT: 83036, 83037 • HbA1c Test Result or Finding <ul style="list-style-type: none"> • CPT CAT II: 3044F, 3046F, 3051F, 3052F <p>Note: Do not include a modifier when using CPT-CAT-II codes</p>
<p>(UOP) Use of Opioids from Multiple Providers ● ●</p> <p>Members 18 years and older, receiving prescription opioids for ≥15 days during the year, who received opioids from multiple providers.</p> <p>Three Rates:</p> <ul style="list-style-type: none"> • Multiple Prescribers – % of members receiving opioids from four or more different prescribers. • Multiple Pharmacies – % of members receiving opioids from four or more different pharmacies. • Multiple Prescribers and Multiple Pharmacies – members receiving opioids from four or more prescribers and four or more pharmacies. <p>Note: A lower rate indicates better performance for all three rates.</p>	<p>Tips</p> <ul style="list-style-type: none"> • Only prescribe opioids when medically necessary, in the lowest effective dose, for the shortest duration necessary. • Identify alternatives to opioids for pain management such as NSAIDS, physical therapy, acupuncture, massage therapy, and corticosteroids when clinically appropriate. • Set expectations early-on regarding controlled-substance prescriptions from other providers and the use of multiple pharmacies 	<p>Opioid Medications</p> <ul style="list-style-type: none"> • Benzhydrocodone • Buprenorphine (transdermal patch and buccal film) • Codeine • Dihydrocodeine • Fentanyl • Hydrocodone • Hydromorphone • Levorphanol • Meperidine • Methadone • Morphine • Opium • Oxycodone • Oxymorphone • Pentazocine • Tapentadol • Tramadol <p>(Opioid medications exclude injectables and opioid-containing cough and cold products).</p>

Lines of Business: Ambetter (Marketplace) = ● Wellcare (Medicare) = ● KanCare (Medicaid) = ●

This document is an informational resource designed to assist licensed healthcare practitioners in caring for their patients. Healthcare practitioners should use their professional judgment in using the information provided. HEDIS[®] measures are not a substitute for the care provided by licensed healthcare practitioners and patients are urged to consult with their healthcare practitioner for appropriate treatment. Codes are subject to change. HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA).

*CPT copyright 2024 American Medical Association (AMA). All rights reserved. CPT is a registered trademark of the AMA.

KDHE-Approved 8/16/2024